

Observations on spacetime symmetry and non-commutativity

Pulak Ranjan Giri*

Theory Division, Saha Institute of Nuclear Physics, 1/AF Bidhannagar, Calcutta 700064, India

T. Shreecharan†

The Institute of Mathematical Science, CIT Campus, Taramani, Chennai 600113, India

(Dated: December 30, 2008)

We consider both the co-ordinates and momenta to be non-commutative and define a non-commutative version of Lorentz symmetry which has a smooth limit to the standard Lorentz symmetry. The Poincaré algebra in this spacetime has also been discussed.

PACS numbers: 03.65.-w, 03.65.Db, 03.65.Ta

The study of non-commutative geometry and the formulation of field theories in such a space-time have become the subject of intense investigation recently [1, 2, 3, 4, 5, 6]. One possible motivation is the appearance of the non-commutative field theory as the limit of certain string theory [7]. However, non-commutativity is the intrinsic nature of quantum theory. For example, the conjugate co-ordinates, (x, p) , of a quantum mechanical model are non-commutative, leading to the famous Heisenberg uncertainty relation

$$\Delta x \Delta p \sim \hbar. \quad (1)$$

In quantum field theory the field operator and its corresponding momentum operator does not commute. It is obvious that in the presence of spacetime non-commutativity the spacetime co-ordinates can not be measured simultaneously, leading to an uncertainty relation. For the co-ordinates x, y being non-commutative one gets an uncertainty relation of the form

$$\Delta x \Delta y \sim \theta. \quad (2)$$

Incorporating spacetime non-commutativity naively in standard quantum mechanics and quantum field theory is fraught with difficulties and subtleties that one needs to tackle.

The introduction of spacetime non-commutativity in general modifies the different symmetries of a system and it may break the symmetries also. So, one of the prime objective is to investigate the spacetime symmetries in non-commutative spacetime. It is known that when the spacetime is non-commutative, with the non-commutative parameter being constant, then the Lorentz symmetry of a system breaks down. However it is possible to restore the Lorentz symmetry if one considers the non-commutative parameter $\theta_{\mu\nu}$ to transform under Lorentz transformation accordingly. However, in 1947 Snyder showed that it is possible to restore Lorentz invariance by a non-commutative spacetime of a definite form [8, 9]. Although, the Poincaré symmetry is usually lost.

In order to handle the problems, associated with the non-commutativity, different forms of non-commutativity has been introduced in literature. For example in [2], the non-commutativity of spacetime deforms the spacetime symmetry algebra of a system. It has been shown that in relativistic case the Lorentz, Poincaré and Conformal algebra is modified. They obtained a one parameter family of conformal generators which form a closed algebra whose commutative limit is smooth. The corresponding non-relativistic counterpart of the symmetry algebra for example Galilean, Schrödinger, non-relativistic conformal algebra are also studied in non-commutative space which leads to a deformed algebra. Another example of getting a consistent Lorentz and Poincaré invariance in non-commutative spacetime is to twist [10, 11] the co-product defined by $\Delta_\theta = \mathcal{T}^{-1} \Delta \mathcal{T}$, where $\mathcal{T} = \exp(-\frac{i}{2} \theta^{\mu\nu} \partial_\mu \otimes \partial_\nu)$. This is known as Drinfel'd twist.

The above mentioned prescription for implementing spacetime symmetries involve only co-ordinate space non-commutativity. However in classical mechanics the phase space coordinates are treated in equal footing. One can consider any set of the phase space variables as the coordinates of the system and the remaining half then becomes the conjugate momenta. Therefore, there is no reason to consider only coordinate non-commutativity and leaving the momenta commutative. In fact it is known that, in quantum mechanics, the generalized momenta P_i in the magnetic field, B , background do not commute, which leads to an uncertainty relation,

$$\Delta P_1 \Delta P_2 \sim B. \quad (3)$$

In this letter we consider both the coordinates and momenta to have deformation. We point out that such a thing is not very exotic and has been introduced and studied previously [5, 12, 13]. We discuss the Lorentz transformation in a phase space where both the co-ordinates and momenta are non-commutative within themselves. Note that in [14] the non-commutative Lorentz symmetry has been obtained by considering only non-commutative space. They show that there exists an invariant length in non-commutative space which re-

mains invariant under Lorentz transformation. It must be pointed out that, what we call as Lorentz transformation is that set of transformations that leave the commutators (4) invariant.

We start our discussion of this letter by taking both the co-ordinates and momenta to be non-commutative as mentioned above. But then the Heisenberg uncertainty principle will be modified in general. So, let us consider the phase space algebra $\mathcal{A}_{\theta,\hat{\theta}}$ due to non-commutativity to be of the form

$$\begin{aligned} [\hat{x}_\mu, \hat{x}_\nu] &= i2\theta_{\mu\nu}, \quad [\hat{p}_\mu, \hat{p}_\nu] = i2\hat{\theta}_{\mu\nu}, \\ [\hat{x}_\mu, \hat{p}_\nu] &= i\hbar(\delta_{\mu\nu} + \theta_\mu^\beta \hat{\theta}_{\nu\beta}). \end{aligned} \quad (4)$$

Here $\theta_{\mu\nu}$ and $\hat{\theta}_{\mu\nu}$ are the non-commutativity parameters for co-ordinates and momenta respectively. Note that the commutator between co-ordinates and its conjugate momenta has been modified, which usually does not happen if one considers only co-ordinates as non-commuting. One can define a modified Planck's constant [12] like $\hbar(\delta_{\mu\nu} + \theta_\mu^\beta \hat{\theta}_{\nu\beta})$. This algebra, $\mathcal{A}_{\theta,\hat{\theta}}$, is not consistent with the standard Lorentz symmetry because, the non-commutativity parameters $\theta, \hat{\theta}$ are considered to be constant and therefore do not transform under the Lorentz transformation. On the other hand the Lorentz algebra is consistent with a phase space where both the co-ordinate space and momentum space are commutative. The algebra $\mathcal{A}$ being

$$[x_\mu, x_\nu] = 0, \quad [p_\mu, p_\nu] = 0, \quad [x_\mu, p_\nu] = i\hbar\delta_{\mu\nu}.$$

Note however that the commutative limit is smooth

$$\lim_{\theta, \hat{\theta} \rightarrow 0} \mathcal{A}_{\theta, \hat{\theta}} = \mathcal{A}. \quad (5)$$

This can be taken as a guideline to get a non-commutative version of Lorentz transformation which would be consistent with the algebra $\mathcal{A}_{\theta,\hat{\theta}}$. In commutative space we know that Lorentz transformation is defined as

$$x'_\mu = \Lambda_\mu^\nu x_\nu, \quad p'_\mu = \Lambda_\mu^\nu p_\nu, \quad (6)$$

which leave the lengths invariant

$$S_x^2 = x'_\mu x'^\mu = x_\mu x^\mu, \quad S_p^2 = p'_\mu p'^\mu = p_\mu p^\mu. \quad (7)$$

Since we want to know how the non-commutative phase space co-ordinates, $\hat{x}_\mu, \hat{p}_\mu$, will transform under the Lorentz transformation (6) we need to know how they are related to the commutative phase space co-ordinates x_μ, p_μ . One possible representation is of the form

$$\hat{x}_\mu = x_\mu - \theta_{\mu\nu} p^\nu, \quad \hat{p}_\mu = p_\mu + \hat{\theta}_{\mu\nu} x^\nu, \quad (8)$$

It is equivalent to the relation

$$x_\mu = \hat{x}_\mu + \theta_{\mu\nu} \hat{p}^\nu, \quad p_\mu = \hat{p}_\mu - \hat{\theta}_{\mu\nu} \hat{x}^\nu, \quad (9)$$

Let us now suppose that under Lorentz transformation the non-commutative phase space co-ordinates transform as

$$\hat{x}'_\mu = x'_\mu - \theta_{\mu\nu} p'^\nu, \quad \hat{p}'_\mu = p'_\mu + \hat{\theta}_{\mu\nu} x'^\nu, \quad (10)$$

which together with (6) and (9) gives the generalization of Lorentz transformation

$$\begin{aligned} \hat{x}'_\mu &= (\Lambda_\mu^\alpha + \theta_{\mu\nu} \Lambda_\beta^\nu \hat{\theta}^{\beta\alpha}) \hat{x}_\alpha + (\Lambda_\mu^\alpha \theta_{\alpha\delta} - \theta_{\mu\nu} \Lambda_\delta^\nu) \hat{p}^\delta, \\ \hat{p}'_\mu &= (\Lambda_\mu^\alpha + \hat{\theta}_{\mu\nu} \Lambda_\beta^\nu \theta^{\beta\alpha}) \hat{p}_\alpha - (\Lambda_\mu^\alpha \hat{\theta}_{\alpha\delta} - \hat{\theta}_{\mu\nu} \Lambda_\delta^\nu) \hat{x}^\delta \end{aligned} \quad (11)$$

One can of course put a restriction

$$\Lambda_\mu^\alpha \theta_{\alpha\delta} = \theta_{\mu\nu} \Lambda_\delta^\nu, \quad \Lambda_\mu^\alpha \hat{\theta}_{\alpha\delta} = \hat{\theta}_{\mu\nu} \Lambda_\delta^\nu, \quad (12)$$

on the transformation (11), which then becomes,

$$\begin{aligned} \hat{x}'_\mu &= (\Lambda_\mu^\alpha + \theta_{\mu\nu} \Lambda_\beta^\nu \hat{\theta}^{\beta\alpha}) \hat{x}_\alpha, \\ \hat{p}'_\mu &= (\Lambda_\mu^\alpha + \hat{\theta}_{\mu\nu} \Lambda_\beta^\nu \theta^{\beta\alpha}) \hat{p}_\alpha. \end{aligned} \quad (13)$$

The restriction (12) is only satisfied in 2-dimensions, but in more than 2-dimensions it is not true in general [2]. Note that when both the non-commutativity parameters are same $\theta = \hat{\theta}$ then the Lorentz transformation for both co-ordinates and momenta are same in 2-dimensions. One can then define a generalized version of Lorentz transformation due to non-commutativity as

$$\widetilde{\Lambda}_\mu^\alpha = \Lambda_\mu^\alpha + \theta_{\mu\nu} \Lambda_\beta^\nu \theta^{\beta\alpha}. \quad (14)$$

We can also consider the Poincaré algebra, $\mathcal{P}$,

$$\begin{aligned} [J^{\alpha\beta}, J^{\gamma\delta}] &= i(-\eta^{\beta\gamma} J^{\alpha\delta} + \eta^{\alpha\gamma} J^{\beta\delta} + \eta^{\delta\alpha} J^{\gamma\beta} - \eta^{\delta\beta} J^{\gamma\alpha}), \\ [p^\alpha, J^{\beta\gamma}] &= i(-\eta^{\alpha\beta} p^\gamma + \eta^{\alpha\gamma} p^\beta), \\ [p^\alpha, p^\beta] &= 0, \end{aligned} \quad (15)$$

where

$$J^{\alpha\beta} = x^\alpha p^\beta - x^\beta p^\alpha. \quad (16)$$

In the context of our non-commutative framework, the operator $\hat{J}^{\alpha\beta}$, obtained replacing the phase space co-ordinates x_μ, p_μ by corresponding non-commutative counter parts $\hat{x}_\mu, \hat{p}_\mu$ in $J^{\alpha\beta}$, becomes

$$\begin{aligned} \hat{J}^{\alpha\beta} &= \hat{x}^\alpha \hat{p}^\beta - \hat{x}^\beta \hat{p}^\alpha \\ &= x^\alpha p^\beta + x^\alpha \hat{\theta}^{\beta\nu} x_\nu - \theta^{\alpha\mu} p_\mu p^\beta \\ &\quad - \theta_\mu^\alpha p^\mu \hat{\theta}^{\beta\nu} x_\nu - < \alpha \rightleftharpoons \beta >, \end{aligned} \quad (17)$$

where $< \alpha \rightleftharpoons \beta >$ refers to all previous terms but with α and β interchanged. Note that the commutators among $\hat{J}^{\mu\nu}$ and $\hat{p}^\mu$ will not close and therefore does not form any algebra. However since we considered (11) as our generalized Lorentz transformation, which is consistent with the standard Lorentz algebra in commutative space (6), the

Poincaré algebra (15) will also be the corresponding algebra in non-commutative space. This is also commented in [14], but there only co-ordinate space non-commutativity is considered.

In conclusion, we discussed the spacetime symmetry transformation for example Lorentz and Poincaré symmetry in non-commutative spacetime. We considered both the co-ordinates and their conjugate to be non-commutative. The reason the spacetime symmetry algebra remains invariant in non-commutative spacetime is that the representation of non-commutative variables like (8) is nothing but going to a new basis.

ACKNOWLEDGMENT

P. R. Giri thanks T. R. Govindarajan for giving him chance to attend the workshop on “Non-commutative geometry and quantum field theory” held on 18 - 24 December, 2008 at IMSc, Chennai, India, where part of this work has been performed.

[†] Electronic address: shreet@imsc.res.in

- [1] M. R. Douglas and N. A. Nekrasov, Rev. Mod. Phys. **73**, 977 (2001).
- [2] R. Banerjee and K. Kumar, Phys. Rev. **D75**, 045008 (2007).
- [3] B. Chakraborty, S. Gangopadhyay and A. Saha, Phys. Rev. **D70**, 107707 (2004).
- [4] T. R. Govindarajan, K. S. Gupta, E. Harikumar, S. Meljanac and D. Meljanac, Phys. Rev. **D77**, 105010 (2008).
- [5] V. P. Nair and A. P. Polychronakos, Phys. Lett. **B505**, 267 (2001).
- [6] A.P. Balachandran, T.R. Govindarajan and B. Ydri, Mod. Phys. Lett. **A15**, 1279 (2000).
- [7] N. Seiberg, E. Witten, JHEP **09**, 032 (1999).
- [8] H. S. Snyder, Phys. Rev. **71**, 38 (1947).
- [9] H. S. Snyder, Phys. Rev. **72**, 68 (1947).
- [10] M. Chaichian, P. P. Kulish, K. Nishijima and A. Tureanu, Phys. Lett. **B604**, 98 (2004).
- [11] M. Chaichian, P. Presnajder, and A. Tureanu, Phys. Rev. Lett. **98**, 151602 (2005).
- [12] O. Bertolami, J. G. Rosa, C. M. L. de Aragao, P. Castorina and D. Zappala, Phys. Rev. **D72**, 025010 (2005).
- [13] P. R. Giri, P. Roy, arXiv:0806.2564v2 [hep-th], Eur. Phys. J. C (in press).
- [14] X. Calmet, Phys. Rev. **D71**, 085012 (2005).

* Electronic address: pulakranjan.giri@saha.ac.in